



Dr. Labovitch's ARTHROSCOPIC KNEE Post-Op

1. Immediate Post-op:

- Keep operative leg Elevated (at least 2 pillows) **as much as possible**
- Ice for the 20 minutes every couple of hours for the first 2-3 days **(DO NOT PLACE ICE DIRECTLY ON THE SKIN. THIS CAN CAUSE SKIN BURNS THAT CAN BE PERMANENT AND CAUSE A WOUND INFECTION)**
- If you are using a cold therapy machine, **DO NOT PLACE THE WRAP DIRECTLY AGAINST THE SKIN. THIS CAN CAUSE SKIN BURNS THAT CAN BE PERMANENT AND CAUSE A WOUND INFECTION**
- **Weight Bearing Status:**
 - ___ Weight bear as tolerated (use crutches only as needed)
 - ___ Partial Weight Bearing (50% of your body weight with crutches)
 - ___ Touch-Down Weight Bearing
 - ___ Non- Weight bearing for ___ day's ___ weeks
 - Other: _____

2. Pain Medication:

- **Expect your pain to be the worst for the first 3 days after surgery**
- If you have been given a **nerve block**, you may be pain free or comfortable for 12-24 hours. When the block begins to wear off, your pain will increase quickly.
- Stay on top of your pain by taking your medication as prescribed
You will be Rx pain medication at your pre-op visit. **IMPORTANT:** Percocet has Tylenol within the medicine. You may take Percocet OR Tylenol for pain. **Do NOT take together**). Please fill prior to your surgery.
- NSAIDs (such as Advil, Motrin, Ibuprofen) may be taken 24 hours after surgery and can be taken with the prescribed pain medication OR Tylenol.
- If prescribed, **Vistaril** can help with nausea, itching, rash, and uncontrolled pain
- Wean off your medication when your pain decreases
- **CONSTIPATION:** Narcotic pain medication can cause constipation. You may use a stool softener (Colace or Dulcolax). Both are sold over-the-counter

3. Dressing/Wound:

- Take down dressing 72 hours after surgery
- Place a band aid or dry dressing over the wounds
- Replace ace bandage for the 1st week after surgery
- **SHOWERS** are allowed 72 hours after surgery
- **NO baths, Jacuzzi, or Sauna for 3 weeks after surgery**
- **Do not** place any gels or ointments on or around the wounds

4. Recommended Activities:

- Begin **Range of Motion** exercises (Flexion and Extension) 2-3 days after surgery
- **Isometric quad exercises:** straight leg raises with the knee extended
- Flex and extend your ankles to increase blood circulation (unless you are in a splint used after foot, leg, or ankle surgery.....or if instructed not to do so by Dr. Labovitch)
- Follow up with Physical Therapy within 1 week after surgery
- Other: _____

5. Nutrition and Sleep:

- It is very important to have a well-balanced diet before and after surgery. This will help with proper wound healing and decrease the chance for complications.
- **DO NOT DRINK ALCOHOL WHILE ON NARCOTIC PAIN MEDICATION.** Please limit your intake of alcohol before and after surgery. This will help with proper wound healing and decrease the chance for complications.
- **DO NOT SMOKE TOBACCO OR ANY OTHER INHALED SUBSTANCES AS THIS IS A KNOWN RISK FACTOR FOR COMPLICATIONS (ESPECIALLY WOUND COMPLICATION, INFECTION, AND DECREASED BONE HEALING)**
- Adequate **sleep** is important. This will help with proper wound healing and decrease the chance for complications.

6. WHAT TO WATCH OUT FOR:

- **PAIN** not controlled by your medication
- Persistent **Drainage** coming out of the wounds
- **Fevers** greater than 100.5 lasting greater than 12 hours

7. Phone numbers and Website:

- **949-520-1012** (during normal business hours)
- **949-520-1012** (after hours)
- **www.ryanslabovitchmd.com** (patient education button)